

Supplementary data from:

Time to diagnosis in Cushing's syndrome: A meta-analysis based on 5367 patients

German Rubinstein¹, Andrea Osswald¹, Eva Hoster², Marco Losa³, Atanaska Elenkova⁴, Sabina Zacharieva⁴, Márcio Carlos Machado⁵, Felicia Alexandra Hanzu⁶, Stephanie Zopp¹, Katrin Ritzel¹, Anna Riester¹, Leah Theresa Braun¹, Ilonka Kreitschmann-Andermahr⁷, Helen L. Storr⁸, Prachi Bansal⁹, María-José Barahona¹⁰, Elisa Cosaro¹¹, Sema Ciftci Dogansen¹², Philip C. Johnston¹³, Ricardo Santos de Oliveira¹⁴, Christian Raftopoulos¹⁵, Carla Scaroni¹⁶, Elena Valassi¹⁷, Steven J.A. van der Werff¹⁸, Jochen Schopohl¹, Felix Beuschlein^{1,19}, Martin Reincke^{1\$}

The associated article has been submitted to the *Journal of Clinical Endocrinology and Metabolism*.

Please cite the original article when using these data.

Figure 1: Mean time to diagnosis of all included studies.

Figure 2: Mean time to diagnosis according to sex (female, male).

Figure 3: Mean time to diagnosis according to age in pituitary CS (<18 years, >=18 years).

Figure 4: Mean time to diagnosis according to year of diagnosis in pituitary CS (before and after 2000).

Figure 1:




Figure 2:




Figure 3:




Figure 4:

